

Evaluation of Nitrogen Fertilization Practices for Surface-Irrigated Lemon Trees - 2008¹

Glenn C. Wright

Department of Plant Sciences, U. of A., Yuma Agriculture Center, Yuma, AZ

Abstract

Lisbon lemons were treated with N levels ranging from 0.5 to 3.0 lbs. N per tree annually. Yield results from the trial show no significant effects of the treatments for the first harvest, but trees treated with the lowest N rate had significantly less yield than trees treated with a higher rate for the second harvest.

Introduction

The University of Arizona first published nitrogen and phosphorous fertilization recommendations for lemons on the sandy soils of the Yuma Mesa in 1961². Using ammonium sulfate for that study, Hilgeman and Rodney reported that for lemon trees on the Superstition sand, 2½ lbs. of N per tree was sufficient for phosphate-fertilized trees up to 4 years of age. They also reported that for those young lemons, there was no improvement in yield as N fertilization increased to 4 lbs. per tree. These authors did not report results for older trees, nor did they correlate their results with lemon leaf nitrogen concentration; instead they noted that when leaf N levels dropped below 1.9% for grapefruit and 2.2% for orange, fertilization was required.

Thirty years later, Doerge *et al.* recommended that N fertilization rates for mature citrus should vary based on leaf concentration as shown in Table 1³. For mature citrus orchards, these authors recommend 2 to 3 lbs. N per tree on sandy soils; however they also suggest higher rates for lemons. It is unclear whether those higher rates refer to 3 lbs. per tree or more.

The most recent work that addressed the lemon fertilization question was done by Sanchez *et al.*, in 2002⁴. In that one-year study, on 8-year old lemon trees, just three rates (0, 1.5 and 3.0 lbs)

Total N in Leaves	Apply this amount of N per tree
%	Lbs.
<2.2	3 - 4
2.2 - 2.3	2 - 3
2.4 - 2.6	1 - 2
2.7 - 2.8	½ - 1
>2.8	0 - ½

Adapted from Doerge *et al.*

¹ The author wishes to thank Mr. Marco Peña and Associated Citrus for their assistance in completing this project. The author would also like to thank the Arizona Citrus Research Council for supporting this research. This is a final report for project 2008-01 – Evaluation of Nitrogen Fertilization Practices for Surface-Irrigated Lemon Trees

² Hilgeman, R.H and D.R. Rodney. 1961. Commercial citrus production in Arizona. University of Arizona Agricultural Experiment Station and Cooperative Extension Service Special Report No. 7. University of Arizona, Tucson, AZ.

³ Doerge, T.A., R.L. Roth and B.R. Gardner. 1991. Nitrogen fertilizer management in Arizona. University of Arizona College of Agriculture Publication No. 191025.

⁴ Sanchez, C.A., G.C. Wright and M. Peralta. 2002. Continued evaluation of N fertilization practices for surface-irrigated lemons. In: G.C. Wright and R. Gibson (eds). 2003 Citrus and deciduous fruit and nut research report. University of Arizona Cooperative Extension Publication No. AZ 1331.

of N per tree were applied via surface irrigation, along with foliar N fertilization and P application. These authors noted that just 1.5 lb. N per tree was sufficient for maximum yield. It should be noted that this study was not conducted on Superstition sand, but rather on Superstition complex soil, a soil in which the sand is topped with a silt layer. Their results are similar to Hilgeman and Rodney's 1961 the recommendation of 1 to 2 lbs. N for lemons planted in Superstition complex. From this survey of the literature, it becomes apparent that there are still no concrete N fertilization recommendations for mature lemons grown in the Superstition sand. Therefore, our objective is to establish and conduct an N fertility trial to establish those recommendations.

Materials and Methods

This experiment was established in a commercial lemon block near County 14th Street and Avenue 3E. There were six treatments, 0.5, 1.0, 1.5, 2.0, 2.5 and 3.0 lbs. (Treatments 1-6) nitrogen per tree. A treatment unit consisted of four adjacent trees in a row, and experimental design was randomized complete block, with seven blocks. Therefore, there were a total of 42 treatment units, (168 trees, 1.54 acres) included in the experiment. Spacing was 20 ft by 20 ft. Each treatment consisted of 7 treatment units, one for each block, with a total of 28 trees, or 0.26 acres. Guard rows separated one treatment from the next where possible. Irrigation is border flood, and normal cultural practices are used.

Nitrogen was applied as UN-32 (32-0-0) or 15-0-0-16S. Most treatments were applied at a rate of 7.5 or 8.0 gallons per acre. For treatments 5 and 6 (2.5 and 3.0 lbs N per acre), a double application was applied as needed. Application details are found in Table 1.

Table 1. Details of N application rates, materials, and timings.

Date	Material	Rate (gpa)	Treatments	Rate	Treatments
31-Jan-08	15-0-0-16S	8.0	1,2,3,4,5,6	15	None
14-Feb-08	32-0-0	7.5	3,4,5,6	15.0	None
28-Feb-08	32-0-0	7.5	4,5,6	15.0	None
11-Mar-08	32-0-0	7.5	1,2,3,4,5,6	15.0	None
1-Apr-08	32-0-0	7.5	5,6	15.0	None
17-Apr-08	32-0-0	7.5	2,3,4	15.0	5,6
1-May-08	15-0-0-16S	8.0	1,2,3,4,5,6		
14-May-08	32-0-0	7.5	2,3,4,5	15.0	6
29-May-08	32-0-0	7.5	4,5,6	15.0	None

Liquid fertilizer was applied to the soil surface using a boom sprayer, one to two days prior to a flood irrigation event.

Initial pick of the experiment was 9-8-08 followed by an additional pick on 10-31-08. In January, 2009, the block was inadvertently harvested, and the final yield data was lost. For each harvest, fruit from each tree was harvested by hand using professional pickers from a local packinghouse. Fruit from each tree was harvested into plastic bins, each holding approximately 800 lbs. From those bins, about 40 lbs of fruit were removed into plastic sample tubs for determination of packout. Fruit from the tubs was optically sorted using a completely automated photographic sorter (Autoline, Inc., Reedley, CA). This sorter is trailer-mounted so that it can be towed to the citrus orchard study site. Each fruit that passes through the sorter was photographed and weighed. Weight, color, exterior quality (% blemish), fruit shape and fruit diameter data was collected for each fruit. Fruit were not physically sorted, but the data collected was stored in a laptop computer that is an integral part of the sorter. Data collected from the sorter were later analyzed and the percentage of fruit from the eight fruit sizes and fruit grades (fancy, choice and juice) as well as fruit peel color and shape were determined. Our results typically show that fruit is larger than is the case if the packout was determined at the packinghouse. This due to the fact that we measure each fruit shortly after it is removed from the trees. In typical packinghouse reports, the percentage of large fruit is less, because fruit shrinks as it moves through the house, and the reports are taken after the fruit is degreened, washed, waxed and dried.

Throughout this experiment, we found no effect of the treatments on fruit grade, shape or peel color, In general, exterior fruit quality ranged from 70 to 75% fancy, about 20% choice, and 5 to 10% juice.

All data was analyzed using SPSS 11.0 for Windows (SPSS Inc., Chicago, Illinois).

Results and Discussion

Partial yields for the 2008-09 season appear in Figure 1. For the 9/8 harvest, yields ranged from 36 lbs. per tree for the trees treated with 2.0 lbs. per tree to 66 lbs. per tree for the trees treated with 0.5 lbs. per tree. Despite the range of yields, we found no significant effect of the treatments upon first harvest yield. For the second harvest on 10/31/08, yields ranged from 233 lbs per tree for the trees treated with 0.5 lbs. per tree, and this value was significantly less than the 281 lbs. per tree recorded for the trees treated with 2.5 lbs. of N annually. We found no effect of the treatments upon fruit packout (Figures 2 and 3).

While it is unfortunate that the final yield was lost, the significant differences noted in the second harvest suggest that the treatments are beginning to have some effect. If so, it is likely that these treatments may have a significant effect upon lemon yield in the 2009/10 harvest season.




Figure 1. 2008-09 season yields for lemons treated with various N levels.




Figure 2. 9/8/08 packout of lemons treated with N levels ranging from 0.5 to 3.0 lbs. of N per tree.




Figure 3. 10/31/08 packout of lemons treated with N levels ranging from 0.5 to 3.0 lbs. of N per tree.